

Quotient Equations and Integrals of Motion for Massive Spinor Field

S.S. Moskaliuk

Bogolyubov Institute for Theoretical Physics
Metrolohichna Str., 14-b, Kyiv-143, Ukraine, UA-03143
e-mail: mss@bitp.kiev.ua

Abstract

In this article a group-theoretical aspect of the method of dimensional reduction is presented. Then, on the base of symmetry analysis of an anisotropic space geometrical description of dimensional reduction of equation for massive spinor field is given. Formula for calculating components of the energy-momentum tensor from the variables of the field factor-equations is derived.

Let us consider massive spinor field in metric

$$ds^2 = dt^2 - \sum_{i=1}^3 A_i^2(t)(dx^i)^2. \quad (1)$$

To this end we shall use the tetrad formalism and present metric in the form

$$ds^2 = g_{\mu\nu}dx^\mu dx^\nu = \eta_{ab} \left(h_\mu^{(a)} dx^\mu \right) \left(h_\nu^{(b)} dx^\nu \right),$$

where $\eta_{ab} = \text{diag}(1, -1, -1, -1)$ is metric tensor in tangent Minkowski space; $h_\mu^{(a)}$ are reference vectors ($a, b = 0, 1, 2, 3$), and here $h_{(a)}^\mu h_{(b)\mu} = \eta_{ab}$, $h_\mu^{(a)} h_{(a)\nu} = g_{\mu\nu}$. It is convenient to choose the reference vectors in this metric in the form of $h_\mu^{(0)} = \delta_\mu^0$, $h_k^{(a)} = \mathcal{A}_k \delta_k^a$. Let us write general covariant Dirac equation

$$(i\gamma^\mu(x)\widehat{\nabla}_\mu - m)\psi(x) = 0. \quad (2)$$

Here $\gamma^\mu(x) = h_{(a)}^\mu \gamma^a$ is 4-vector in respect to general coordinate transformations;

$$\widehat{\nabla}_\mu = \partial_\mu + \frac{1}{4} C_{abc} h_\mu^{(c)} \gamma^b \gamma^a$$

is spinor covariant derivative; C_{abc} are Ricci rotation coefficients.

For constant γ -matrices we use the representation

$$\gamma^0 = \begin{pmatrix} \sigma^0 & 0 \\ 0 & -\sigma^0 \end{pmatrix}, \quad \gamma^k = \begin{pmatrix} 0 & \sigma^k \\ -\sigma^k & 0 \end{pmatrix}, \quad \gamma^5 = \begin{pmatrix} 0 & \sigma^0 \\ \sigma^0 & 0 \end{pmatrix},$$

where σ^μ are Pauli matrices.

Christoffel symbols and Ricci rotation coefficients $\Gamma_{kk}^0 = \mathcal{A}_k \dot{\mathcal{A}}_k$, $\Gamma_{k0}^k = \dot{\mathcal{A}}_k / \mathcal{A}_k$, $C_{k0k} = \dot{\mathcal{A}}_k / \mathcal{A}_k$ (no summation over k) differ from zero. The spinor covariant derivatives are $\widehat{\nabla}_0 = \partial/\partial t$, $\widehat{\nabla}_k = \partial/\partial x^k + \dot{\mathcal{A}}_k \gamma^0 \gamma^k / 2$. Then the equation (2) acquires the form

$$\left(i \frac{\partial}{\partial t} + \frac{1}{2} i \sum_k \frac{\dot{\mathcal{A}}_k}{\mathcal{A}_k} + i \frac{\alpha^k}{\mathcal{A}_k} \frac{\partial}{\partial x^k} - \beta m \right) \psi(x),$$

$$\alpha^k = \gamma^0 \gamma^k, \quad \beta = \gamma^0.$$

The second summand can be easily eliminated, substituting $\psi(x)$ in the form $\psi(x) = (\mathcal{A}_1 \mathcal{A}_2 \mathcal{A}_3)^{-1/2} \Psi(x)$. As a result we obtain

$$\left(i \frac{\partial}{\partial t} + i \frac{\alpha^k}{\mathcal{A}_k} \frac{\partial}{\partial x^k} - \beta m \right) \Psi(x) = 0. \quad (3)$$

Due to the homogeneity of metric the bispinor $\Psi_{kr}(x) (r = \pm 1)$ can be presented in the form

$$\Psi_{kr}(x) = e^{i\mathbf{k}\mathbf{x}} X_{kr}(t), \quad X_{kr}(t) = \begin{pmatrix} f_{kr}(t) \\ \phi_{kr}(t) \end{pmatrix}. \quad (4)$$

Here spinors $f_{kr}(t)$, $\phi_{kr}(t)$ depend only on time; index r defines the projection of spin on the direction of movement.

Substituting (4) in (3), we get a set of equations for spinors $f_{kr}(t)$ and $\phi_{kr}(t)$:

$$\begin{aligned} \frac{df_{kr}}{dt} &= -i (m f_{kr} + r q_s \sigma^s \phi_{kr}), \\ \frac{d\phi_{kr}}{dt} &= -i (-m \phi_{kr} + r q_s \sigma^s f_{kr}), \\ (s &= 1, 2, 3), \end{aligned} \quad (5)$$

where $q_k = k_k / \mathcal{A}_k$ are components of physical momentum of particle.

In spherical coordinates we can write

$$\begin{aligned} q_1 &= \frac{k_1}{a\alpha_1} = \frac{k \sin(\theta) \cos(\varphi)}{a\alpha_1}, \quad q_2 = \frac{k_2}{a\alpha_2} = \frac{k \sin(\theta) \sin(\varphi)}{a\alpha_2}, \\ q_3 &= \frac{k_3}{a\alpha_3} = \frac{k \cos(\theta)}{a\alpha_3}, \quad q^2 = q_1^2 + q_2^2 + q_3^2 = \frac{k^2}{a^2} \mu^2 = \frac{k^2}{g^2}, \quad \omega^2 = \frac{k^2}{g^2}. \end{aligned}$$

To find a solution of the set (5) we proceed as follows. Let us choose orthogonal spinors R_{1r} and R_{2r} , describing the independent chiral states in the massless case in the form of

$$R_{1r} = \begin{pmatrix} \sqrt{\frac{q-rq_3}{q}} e^{-i\Phi/2} \\ -r \sqrt{\frac{q+rq_3}{q}} e^{i\Phi/2} \end{pmatrix},$$

$$R_{2r} = \begin{pmatrix} r \sqrt{\frac{q+rq_3}{q}} e^{-i\Phi/2} \\ \sqrt{\frac{q-rq_3}{q}} e^{i\Phi/2} \end{pmatrix},$$

$$\begin{aligned} R_{ir}^+ R_{is} &= 2\delta_{rs}, & R_{1r}^+ R_{2s} &= 2s, \\ R_{2r}^+ R_{1s} &= 2r, & r &\neq s, \\ R_{2r} &= -i\sigma_2 \overset{*}{R}_{1r}, & R_{1r} &= i\sigma_2 \overset{*}{R}_{2r}, \end{aligned}$$

where $\tan(\Phi) = \frac{\alpha_1}{\alpha_2} \tan(\phi)$.

From these spinors we can construct new spinors

$$\begin{aligned} P_{1r} &= \frac{q}{\omega} R_{1r} + \frac{m}{\omega} R_{2r}, \\ P_{2r} &= \frac{q}{\omega} R_{2r} - \frac{m}{\omega} R_{1r}, & P_{3r} &= -i\sigma_2 \overset{*}{P}_{2r}, \\ P_{4r} &= -i\sigma_2 \overset{*}{P}_{1r}, & P_{ir}^+ P_{ir} &= 2, \\ P_{1r}^+ P_{3r} &= -P_{2r}^+ P_{4r} = 4mq/\omega^2. \end{aligned} \tag{6}$$

In massless theory the spinors P_{1r} and P_{2r} describe independent chiral states. If $m \neq 0$, then the transitions $f_{kr} \leftrightarrow \phi_{kr}$ take place with amplitudes proportional to m/ω .

Solution of the set of equations (5) can be found in the form of

$$\begin{aligned} f_{kr}(t) &= \frac{1}{2} \omega^{-1/2} \left[\sqrt{\omega - m} P_{1r} \overset{*}{\alpha}_{kr} e_+ + r \sqrt{\omega + m} P_{3r} \beta_{kr} e_- \right], \\ \phi_{kr}(t) &= \frac{1}{2} \omega^{-1/2} \left[r \sqrt{\omega + m} P_{2r} \overset{*}{\alpha}_{kr} e_+ + \sqrt{\omega - m} P_{4r} \beta_{kr} e_- \right], \\ e_{\pm} &= \exp \left(\pm i \int_{t_0}^t K_0(t') dt' \right). \end{aligned} \tag{7}$$

Hence,

$$\begin{aligned} \overset{*}{\alpha}_{kr}(t) &= \frac{1}{2} \omega^{-1/2} e_- \left[\sqrt{\omega - m} P_{1r}^+ f_{kr} - r \sqrt{\omega + m} P_{2r}^+ \phi_{kr} \right], \\ \beta_{kr}(t) &= \frac{1}{2} \omega^{-1/2} e_+ \left[r \sqrt{\omega + m} P_{3r}^+ f_{kr} + \sqrt{\omega - m} P_{4r}^+ \phi_{kr} \right]. \end{aligned} \tag{8}$$

Using the straightforward substitution and taking into account the normalization conditions, one can verify that $|\alpha_{kr}|^2 + |\beta_{kr}|^2 = 1$. Differentiating (8) by time, we come to a set of linear differential equations of the first order for $\alpha_{kr}(t)$ and $\beta_{kr}(t)$:

$$\begin{aligned}\dot{\alpha}_{kr}^* &= \left(\frac{w}{2} + i\frac{\hat{w}_\perp}{2}\right) \beta_{kr} e_-^2 - i\frac{w_3}{2} \alpha_{kr}^*, \\ \dot{\beta}_{kr} &= \left(-\frac{w}{2} + i\frac{\hat{w}_\perp}{2}\right) \alpha_{kr}^* e_+^2 + i\frac{w_3}{2} \beta_{kr}, \\ q_\perp^2 &= q_1^2 + q_2^2,\end{aligned}\tag{9}$$

with

$$\begin{aligned}w &= \frac{q_3 \dot{q} - \dot{q}_3 q}{\omega q_\perp} \left(\frac{q^2}{\omega^2} - \frac{m^2}{\omega^2}\right) + 2\frac{m^2 \dot{w}}{\omega^3}; \quad \hat{w}_\perp = r\frac{q_\perp}{\omega} \dot{\Phi}; \\ \hat{w}_3 &= r\frac{q_3}{\omega} \dot{\Phi} \left(\frac{q^2}{\omega^2} - \frac{m^2}{\omega^2}\right) + 2r\frac{m^2 q_\perp}{\omega^3} \dot{\Phi}.\end{aligned}\tag{10}$$

In practice it is more convenient to go from two complex-valued functions $\alpha_{kr}(t)$ and $\beta_{kr}(t)$ to three real-valued functions $S_{kr}(t) = |\beta_{kr}^2|^2$, $U_{kr}(t) = -2\text{Re}(\alpha_{kr}\beta_{kr}^* e_-^2)$, $V_{kr}(t) = -2\text{Im}(\alpha_{kr}\beta_{kr}^* e_-^2)$, for which it is possible to derive a set of three linear equations

$$\begin{aligned}\dot{S}_{kr} &= \frac{w}{2} U_{kr} - \frac{\hat{w}_\perp}{2} V_{kr}, \\ \dot{U}_{kr} &= w(-2S_{kr} + 1) + (-\hat{w}_3 + 2K_0) V_{kr}, \\ \dot{V}_{kr} &= -\hat{w}_\perp(-2S_{kr} + 1) - (-\hat{w}_3 + 2K_0) U_{kr}\end{aligned}\tag{11}$$

with initial conditions $S_{kr} = U_{kr} = V_{kr} = 0$ at $t = t_0$.

As it can be seen, the structure of the set of equations is similar to that of the set of equations, derived for scalar field [1] and vector massless field [2].

Using the functions and notation introduced above, we evaluate the vacuum averages of normally ordered operator of energy-momentum tensor $T_{\mu\nu}(t) = \langle 0_{in} | N_t T_{\mu\nu}(t, \mathbf{x}) | 0_{in} \rangle$ for spinor field [2]:

$$\begin{aligned}
T_0^0(t) &= \frac{1}{(2\pi)^3 a^4} \sum_r \int d^3 \mathbf{k} \omega S_{kr}, \\
T_1^1(t) &= \frac{1}{(2\pi)^3 a^4} \sum_r \int d^3 \mathbf{k} q_\perp^2 (\mathcal{X}_{kr} + \mathcal{X}_{kr} \cos(2\Phi) - \mathcal{Z}_{kr} \sin(2\Phi)), \\
T_2^2(t) &= \frac{1}{(2\pi)^3 a^4} \sum_r \int d^3 \mathbf{k} q_\perp^2 (\mathcal{X}_{kr} - \mathcal{X}_{kr} \cos(2\Phi) + \mathcal{Z}_{kr} \sin(2\Phi)), \\
T_3^3(t) &= \frac{1}{(2\pi)^3 a^4} \sum_r \int d^3 \mathbf{k} q_3^2 \mathcal{Y}_{kr}, \quad T_i^0(t) = 0, \quad i = 1, 2, 3, \\
T_2^1(t) &= \frac{1}{(2\pi)^3 a^4} \sum_r \int d^3 \mathbf{k} q_\perp^3 \mathcal{X}_{kr} \sin(2\Phi), \\
T_3^1(t) &= \frac{1}{(2\pi)^3 a^4} \sum_r \int d^3 \mathbf{k} q_\perp q_3 [(\mathcal{X}_{kr} + \mathcal{Y}_{kr}) \cos(\Phi) - \mathcal{Z}_{kr} \sin(\Phi)], \\
T_3^2(t) &= \frac{1}{(2\pi)^3 a^4} \sum_r \int d^3 \mathbf{k} q_\perp q_3 [(\mathcal{X}_{kr} + \mathcal{Y}_{kr}) \sin(\Phi) - \mathcal{Z}_{kr} \cos(\Phi)]. \quad (12)
\end{aligned}$$

Here the following notation is used:

$$\begin{aligned}
\mathcal{X}_{kr}(S_{kr}, U_{kr}) &= \frac{S_{kr}}{\omega} + \frac{1}{2} \frac{q_3 U_{kr}}{\omega q_\perp} \left(\frac{q^2}{\omega^2} - \frac{m^2}{\omega^2} \right) + \frac{m^2}{\omega^3} U_{kr}, \\
\mathcal{Y}_{kr}(S_{kr}, U_{kr}) &= \frac{S_{kr}}{\omega} - \frac{1}{2} \frac{q_\perp U_{kr}}{\omega q_3} \left(\frac{q^2}{\omega^2} - \frac{m^2}{\omega^2} \right) + \frac{m^2}{\omega^3} U_{kr}, \\
\mathcal{Z}_{kr}(V_{kr}) &= -\frac{1}{2} \frac{V_{kr}}{q_\perp}.
\end{aligned}$$

Consequently, we again have reduced analysis of energy-momentum tensor of quantized massive spinor field to analysis of the set of equations for functions S_{kr} , U_{kr} , V_{kr} .

References

- [1] Moskaliuk S.S. Quotient equations and integrals of motion for scalar field // Hadronic Journal. – 2001. – **24**, No. 5. – P. 531–534.
- [2] Moskaliuk S.S. Quotient equations and integrals of motion for vector massless field // Hadronic Journal. – 2001. – **24**, No. 4. – P. 435–448.